

Your smear test results

This leaflet tells you about your results.





This leaflet provides information about the results of your smear

test, how they will be monitored and any treatment you may need.

If you feel anxious about your test results, you may find it helpful to talk about it with someone. Your nurse or GP can give you further information, answer your questions and reassure you that changes are common and can be easily and effectively treated.

It's important to continue to go for screening every time you're invited.

Your risk of cervical cancer increases if:

- you are or have ever been sexually active this includes penetrative sex and other types of sexual activity, such as skin-to-skin contact of the genital area, or using sex toys
- you smoke tobacco smoke affects the cells in your cervix.

It saves around 5,000 lives every year in the UK and prevents 8 out of 10 cervical cancers from developing.

It can pick up changes to your cells even if you look and feel healthy and have no symptoms.

don't ignore the benefits of getting screened

lt can stop cervical cancer before it starts.

It's your best protection against cervical cancer even if you've been immunised against HPV.

Your results

- Smear tests are designed to pick up any changes so they can be easily and effectively treated and monitored.
- Changes in cells are very common.
- Any treatment you may need is usually simple and you can almost certainly be treated as an outpatient.

Make an appointment with your GP as soon as possible if you have unusual discharge, or bleeding after sex, between periods or after the menopause.

Your results letter will tell you if there are:

- no changes to your cells
- not enough cells to test
- minor changes to your cells
- changes which require further investigation.

No changes to your cells

- In most cases your letter will say 'no changes'.
- You'll be invited to have another smear test in three or five years as usual.

Minor changes to your cells

- These sometimes clear up on their own and do not require any treatment.
- These changes will be monitored to check that they have cleared up so you'll be invited to have another smear test in a few months' time.

Not enough cells to test

- Sometimes there are not enough cells in the sample to examine.
- This is not unusual and you'll be invited to repeat the test in three months' time or to have further investigations at a colposcopy clinic (see page 6).

Changes which require further investigation

 If changes are detected which require further investigation you'll be referred to a colposcopy clinic (see page 6) for examination to check whether the changes need to be monitored or treated.

Your questions answered

What happens at the colposcopy clinic?

A colposcopy is an examination of the cervix using a special microscope called a colposcope. The colposcope looks like a large pair of binoculars on a stand. It doesn't go inside you. The specialist will gently insert a speculum into your vagina and will look at your cervix through the colposcope. The colposcope magnifies your cervix so that the specialist can see where the changes are and what they look like. Detailed information about colposcopy will be sent to you with your appointment for the colposcopy clinic.

What about treatment?

Sometimes treatment isn't necessary. If this is the case, the specialist will explain why and will arrange for you to have smear tests more often. In some cases you'll be asked to come back to the clinic for further colposcopy examinations.

If necessary you can have a local anaesthetic and you should feel no more than slight discomfort. The specialist treating you will explain what to expect.

Your follow-up appointment will be with your specialist, nurse or GP.

You can bring a friend or relative to your appointment.

What causes changes in my cervix?

Most changes in the cells of the cervix are caused by a virus called the human papilloma virus (HPV). HPV is very common – 8 out of 10 people in Scotland will catch it at some point in their lives. Because HPV usually has no symptoms, many people have it for months or years without knowing it.

Your body fights off most HPV infections naturally, but about 1 in 10 infections are harder to get rid of.

The types of HPV that can cause changes in the cells of your cervix are transmitted through sexual contact. This includes penetrative sex as well as other types of sexual activity, such as skin-to-skin contact of the genital area, or using sex toys.

Can the test detect all changes?

No. The test can sometimes miss changes and changes can also happen between tests, so it's important to go for a smear test every time you're invited.

Are there any symptoms?

There are usually no symptoms with changes in cervical cells and sometimes there are no symptoms with early-stage cervical cancer. Make an appointment with your GP as soon as possible if you have unusual discharge, or bleeding after sex, between periods or after the menopause.

Want to know more?

Talk to your nurse or GP, visit nhsinform.scot or call 0800 22 44 88 (textphone 18001 0800 22 44 88). The helpline is open every day 8 am to 10 pm and also provides an interpreting service.

Jo's Cervical Cancer Trust: visit jostrust.org.uk or call 0808 802 8000.

This resource may also be made available on request in the following formats:





🕻 0131 314 5300

nhs.healthscotland-alternativeformats@nhs.net

Published by NHS Health Scotland

1 South Gyle Crescent Edinburgh EH12 9EB © NHS Health Scotland 2017

First published 2016

All rights reserved. Material contained in this publication may not be reproduced in whole or part without prior permission of NHS Health Scotland (or other copyright owners). While every effort is made to ensure that the information given here is accurate, no legal responsibility is accepted for any errors, omissions or misleading statements.

NHS Health Scotland is a WHO Collaborating Centre for Health Promotion and Public Health Development.

www.healthscotland.scot